

# EGFR S768\_V769delinsIL(S768I&V769L)/BaF3 CBP73181

## Contents

I. Introduction.....	1
II. Background.....	1
III. Representative Data.....	1
1. WB of EGFR S768_V769delinsIL/BaF3 .....	2
2. Anti-proliferation assay .....	2
IV. Handling Procedure for Flask Cultures.....	2
V. Subculturing Procedure.....	3
VI. Cryopreservation Procedure.....	3
VII. Performing the Anti-proliferation Assay.....	3
1. Materials to Be Supplied by the User.....	3
2. Protocol for the Cell Viability Assay.....	4
VIII. References.....	5



# EGFR S768\_V769delinsIL(S768I&V769L)/BaF3 CBP73181

## I. Introduction

Cell Line Name: EGFR S768\_V769delinsIL(S768I&V769L)/BaF3

Host Cell: Ba/F3

Stability: 16 passages(in-house test, that not means the cell line will be instable beyond the passages we tested.)

Application: Useful for the study of enzyme kinetics, screening inhibitors, and selectivity profiling.

Freeze Medium: 90% FBS+10% DMSO

Complete Culture Medium: RPMI-1640+10%FBS+2ug/ml puromycin

Mycoplasma Status: Negative


## II. Background

EGFR is the receptor for members of the EGF family and is a transmembrane glycoprotein that has tyrosine kinase activity. EGFR S768\_V769delinsIL results in a deletion of two amino acids in the protein kinase domain of the Egfr protein from aa 768 to aa 769, combined with the insertion of an isoleucine (I) and a leucine (L) at the same site (UniProt.org). S768\_V769delinsIL has been identified in the scientific literature (PMID: 17045698, PMID: 27304188), but has not been biochemically characterized and therefore, its effect on Egfr protein function is unknown (PubMed, Jul 2019).

## III. Representative Data



## 1. WB of EGFR S768\_V769delinsIL(S768I&V769L)/BaF3



## 2. Anti-proliferation assay

### IV. Handling Procedure for Flask Cultures

The flask was seeded with cells grown and completely filled with medium at Cobioer.

1. Upon receipt visually examine the culture for macroscopic evidence of any microbial contamination.

Using an inverted microscope (preferably equipped with phasecontrast optics), carefully check for any evidence of microbial contamination.

2. Incubate the flask in an upright position for several hours at 37°C.

After the temperature has equilibrated, aseptically remove the entire contents of the flask and centrifuge at 1000 rpm for 5 minutes. Remove shipping medium and save for reuse. Resuspend the cell pellet in 10 ml of this medium.

3. From this cell suspension remove a sample for a cell count and viability. Adjust the cell density of the suspension to  $3 \times 10^5$  viable cells/ml in the shipping medium.



4. Incubate the culture, horizontally at 37°C in a 5% CO<sub>2</sub> in air atmosphere. Maintain the cell density of the culture as suggested under the subculture procedure.

## V. Subculturing Procedure

Cultures can be maintained by the addition of fresh medium or replacement of medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at  $3 \times 10^5$  viable cells/mL. Do not allow the cell density to exceed  $2 \times 10^6$  cells/mL.

Interval: Maintain cultures at a cell concentration between  $3 \times 10^5$  and  $2 \times 10^6$  viable cells/mL. Medium Renewal: Add fresh medium every 2 to 3 days (depending on cell density)

## VI. Cryopreservation Procedure

1. Transfer cell suspension ( $4-8 \times 10^6$  cells) to a centrifuge tube and spin at approximately 1000rpm for 5 minutes.
2. Discard supernatant and resuspend cells in cryopreservation medium (90% FBS+10%DMSO).
3. Transfer the cells into Freezing Tube, 1ml/vial.
4. Place Freezing Tube in a Frosty container that is kept at room temperature and has sufficient isopropanol.
5. Place the Frosty container into the -80°C freezer overnight.
6. On next day, transfer the vials into liquid nitrogen.

## VII. Performing the Anti-proliferation Assay

### 1. Materials to Be Supplied by the User

- Compounds to be tested



- RPMI-1640 (Gibco #C11875500CP)
- FBS (Gibco #10099-141C )
- DMSO
- CellTiter-Glo® Luminescent Cell Viability Assay (Promega, Cat.No.: G7573)
- 96 Well Assay Plate (White Plate, Clear Bottom with Lid Tissue Culture Treated Polystyrene 1/Pack, Corning #3610)
- T25 Flask
- 96 Well Storage Microplate( Corning#3357)
- Synergy H1 Hybrid Multi-Mode Reader(Biotek)

## 2. Protocol for the Anti-proliferation Assay

- 1) Take logarithmic growth cells, centrifuge and discard the culture supernatant, resuspend the centrifuged cells in fresh RPMI-1640+10%FBS medium.
- 2) Inoculate the resuspended cells into a 96-well cell culture plate with a white wall and transparent bottom, 100ul/well of cell suspension, inoculate two culture plates, and place in a 37°C cell incubator for overnight culture.
- 3) The next day, take one of the 96-well plates inoculated with cells, add 100ul/well cell titer glo detection reagent and leave it for 60 minutes, read the value, and define it as G0 data.
- 4) Take another parallel plate, and take 11.1 ul/well of compound from the previously diluted 10\*compound concentration dilution plate to this plate , And set up another DMSO control well, continue to incubate in a 37°C cell incubator for 72 hours.
- 5) Take out the 96-well plate that has been treated with the compound for 72 hours from the incubator, add 100ul/well cell titer glo detection reagent and leave it for 60 minutes, read the value, and define it as G3



data.

6) Calculate the cell proliferation rate corresponding to each well according to the following formula %Proliferation = (Test compound well G3-G0 average value) / (DMSO control well G3 average value-G0 average value)\*100.

7) According to the corresponding proliferation rate and concentration of each gradient concentration hole, use Prism Graphpad5.0 software to fit the gradient curve of cell proliferation, and calculate the GI50 of the compound (GI50 is defined as the corresponding compound concentration when the cell proliferation rate is 50% , The fitting formula in the software is as follows:  $Y=Bottom + (Top-Bottom)/(1+10^{((LogIC50-X)*HillSlope)})$ ).

## VIII. References

1. Clinicopathologic features of non-small-cell lung cancer with EML4-ALK fusion gene. (PMID: 20183914) Takahashi T ... Date H *Annals of surgical oncology* 2010 3 22 40
2. KRAS mutation status in primary nonsmall cell lung cancer and matched metastases. (PMID: 20336783) Cortot AB ... Hainaut P *Cancer* 2010 3 22 40
3. Are there imaging characteristics associated with epidermal growth factor receptor and KRAS mutations in patients with adenocarcinoma of the lung with bronchioloalveolar features? (PMID: 20087229) Glynn C ... Ginsberg MS *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer* 2010 3 22 40
4. TP53, EGFR, and KRAS mutations in relation to VHL inactivation and lifestyle risk factors in renal-cell carcinoma from central and eastern



Europe. (PMID: 20137853) Szymańska K ... Brennan P *Cancer letters*  
2010 3 22 40

5. Genetic and proteomic features associated with survival after  
treatment with erlotinib in first-line therapy of non-small cell lung cancer  
in Eastern Cooperative Oncology Group 3503. (PMID:  
20035238) Amann JM ... Carbone DP *Journal of thoracic oncology :*  
*official publication of the International Association for the Study of Lung*  
*Cancer* 2010 3 22 40

